Reproductive Health and Preventing Pregnancy

Santana Eastman, RN

Gale Johnsen, PhD, APRN

Spring 2018

IN 2017:

39% of High School students

in Maine said they were

having sex.

DECIDING TO BECOME SEXUALLY ACTIVE?

- · 3% of Maine teens have sex before the age of 13
- 39% have sex by the time they graduate high school
- · Why do we decide to become sexually active?
- - We think it is expected.
- Wanting to be cared about.
- · We care about the other person.
- Other thoughts?

TRENDSINSEXUAL BEHAVIORS 1991-2015

- Percentages decreased since 1991:
- -Those having sexual intercourse
- -Those having sexual intercourse before age 13
- -Drank alcohol or used drugs before last intercourse
- There has been increased use of condoms and birth control.





MAINE INTEGRATED YOUTH HEALTH SURVEY 2017 & NATIONAL YOUTH RISKY BEHAVIOR SURVEY 2015

Let's Look at the

Numbers

21% of teens drank alcohol or used drugs before last sexual encounter

43% of teens did not use a condom during last sexual intercourse

73% did not use birth control pills, IUD or implant; shot, patch or birth control ring during last sexual encounter

3.9% of teens had sexual intercourse before age 13

30.1% of your teen peers are sexually active with at least one person in the last 3 months

10.2% of sexually active teens across the country were ever test for HIV

HAVING DIFFICULT CONVERSATIONS....

- I don't want to become sexually active with you!
- Just because | have been in the past doesn't mean | want to continue!
- Do you have an STD?
- | want you to wear a condom.
- Mom... just had sex (think)

VIOLENCE AND SEXUAL ACTIVITY 2015

-10% of adolescents are hit, slapped, or physically hurt on purpose by their partner in 2015 during sexual activity.

-18% of girls and 11% of boys "forced" to have sexual contact of any type during their lifetimes.

-8% have been forced into sexual activity they did not want to participate in - 6% males, 10% females

GENDERANDSEXUALIDENTITY

- -Why do we ask everyone at the Health Center what their gender identity is and their sexual orientation?
- -3% of teens identify as transgender
- -2% gay
- -2% lesbian
- -3% unsure or gender non-conforming
- -Important because needs and health concerns are different

WHATARE THE CHANCES I'LL BE COME PREGNANT...

• If a female teen has intercourse without

contraception for one year, there is a 90%

chance she will become pregnant.

• 77% of teen pregnancies are unintended.

WHATARE THE CHANCES OF CONTRACTING STI...

· One out of four sexually active teens

develop an STI within one year

TEEN PREGNANCY IN MAINE

- Maine ranked 5th LOWEST birth rate for young women 15-19 in 2015.
- Our birthrate has dropped 52% since 1991. This is a result of education and access to reproductive health care services.
- 93% were to non-married females.

POP QUIZ





to raise a child from birth to 17 years old (not including

college savings)

Not including the annual cost

of college!

\$45,370 Private

\$20,090

Public

*Includes room and board.

The Cost of Raising a Child:

Where does the money go?

Food

Housing

Transportation

Health Care

6% Clothing 16%

Child Care

& Education

Misc.

Annual Housing Cost per Child



VS

Rural \$2,400 **Annual Food Costs by Age**



Cost by Region



Rural Areas \$193,020

*U.S. average for middle-income, married-couple families. Total costs are from birth through age 17.

Source: Expenditures on Children by Families, 2015.

\$233,610

BIRTH CONTROLAND MAINE TEENS

- 16% no birth control with first sex
- 8% don't use ANY protection.
- 18% report

 drugs/etoh when

 having sex, males more
 than females
- 60% using condoms

49% of females who are sexually active are using the pill or a hormone contraception, but still not using condoms consistently - only 18%

HISTORYOFBIRTH CONTROL

- Ancient Egypt sponge
- Ancient Greece spermicides
- England 15th century chastity belts
- Europe 16th century condoms from skins
- USA vulcanizing rubber-condoms 1839

Europe/USA-1930's - cervical cap/diagphrams

USA-1957 - first pill to control menses


Marketed 1960's as birth control pills











TOBECOME PREGNANT:

Have sex.

And don't use birth control.

TONOTBECOME PREGNANT:

Don't have sex.

Or use birth control.

Abstinence

A behavior that prevents pregnancy by not allowing sperm into (or near) the vagina

Prevents sexually transmitted infection

Safe, easy, and convenient

Free, and 100% effective!

People are abstinent for many reasons, including to prevent pregnancy.

Withdrawal (Pull Out Method)

Withdrawal is also called coitus interruptus or the "pull out method"

Something you can do during vaginal intercourse

to prevent pregnancy

Free, safe, easy, and convenient

75% effective

Condoms

Worn on the penis

Made of latex or plastic

Prevent pregnancy and sexually transmitted infection

Can be used for vaginal, anal, or oral sex

Safe, effective, and easy to get

Cost about \$1 each, but are sometimes available for free

75% effective



Birth Control Vaginal Ring (Nuva Ring)

A small ring you put in your vagina once a month for three weeks to prevent pregnancy Safe, effective, and convenient Easy to get with a prescription Costs about \$15-\$70 a month 99% effective



Ortho Evra "The Patch"

- ✓ Small patch that sticks to your skin
- ✓ Safe, effective, and convenient
- ✓ Easy to get with a prescription
- ✓ Costs \$15-\$70/month
- ✓ 99% Effective





0.15 mg-0.02 mg/24 hrs

Ortho Evra®

(norelgestromin/ethinyl estradiol transdermal system)

Birth Control Pills

Take a pill each day to prevent pregnancy
Safe, effective, and convenient Easy to get with a prescription
Cost about \$5-\$50 each month
99% effective



Implant "The Rod"

- ✓99.9 % Effective
- ✓ Hormone Based
- ✓Inserted in the Upper Arm
- ✓ Lasts for 3 years
- ✓In office Procedure:
 - ✓ Insertion
 - ✓ Removal



Birth Control Shot (Depo-Provera)

A shot that prevents pregnancy
Safe, effective, and convenient
Easy to get with a prescription
Lasts for three months
Costs \$35-\$75

99% effective



IUD Intrauterine Device

Small, "T-shaped" device inserted into the uterus to prevent pregnancy


Safe, effective, and long lasting

Must be inserted by a health care provider

Costs between \$500 and \$1,000 up front, but lasts up to 12

years

99% effective



Breastfeeding as Birth Control

Sometimes called LAM (Lactational Amenorrhea Method)

A natural way to prevent pregnancy after giving birth

Effective, safe, convenient, and free Lasts for up to six months after giving birth Less than 75% effective

Emergency Contraception

Birth control you can use to prevent pregnancy up
to five days
after unprotected sex
Safe and effective
Available at health centers and drugstores
Costs vary from \$10 to \$70

NDC 51285-942-88


PlanB One-Step

(levonorgestrel) tablet, 1.5 mg Emergency Contraceptive

Reduces the chance of pregnancy after unprotected sex (ie, if a regular birth control method fails or after sex without birth control)

Not for regular birth control.

Rx only for women younger than age 17



Take as soon as possible within 72 hours (3 days) after unprotected sex. The sooner you take it, the better Plan B[®] One-Step will work.

1 Tablet Levonorgestrel 1.5mg

STERILIZATION

• Tubal Ligation

•Vasectomy

• Should be thought of us a permanent decision.